



*With Compliments
of the Author*

Zagreb, Croatia

Volume 15, Number 1, March 2021

G. MacDonald¹, L. W. Marcoux¹, M. Mastnak¹, M. Omladič² and H. Radjavi

*A note on the structure of
matrix *-subalgebras with scalar diagonals*

OPERATORS AND MATRICES

AIMS AND SCOPE

'Operators and Matrices' (OaM), aims towards developing a high standard international journal which will publish top quality research papers in matrix and operator theory and their applications. The journal will publish mainly pure mathematics, but occasionally papers of a more applied nature could be accepted providing they have a nontrivial mathematical content. OaM will also publish expository papers and relevant book reviews.

OaM is published quarterly, in March, June, September and December.

SUBMISSION

Manuscripts should be submitted through OaM page: oam.ele-math.com, by e-mail of Editorial office, or directly to any one of the editors working in a related subject area.

Manuscripts are accepted for refereeing on the understanding that the same work has not been published (except in the form of an abstract), and is not under consideration for publication elsewhere.

Authors are advised to send notification of submission by e-mail to oam@element.hr (author(s), title, number of pages, name of editor to whom paper was sent to).

In order to facilitate refereeing, copies of those papers (whether by the author or someone else) which are essential and referred to in the manuscript but are not conveniently accessible, should be enclosed.

The publisher strongly encourages submission of manuscripts written in TeX or one of its variants LaTeX, AMSTeX or AMSLaTeX. On acceptance of the paper, authors will be asked to send the file by electronic mail. The file must be the one from which the accompanying manuscript (finalized version) was printed out. A postscript or pdf file of this final version must also be enclosed.

TITLE PAGE

The following data should be included on title page: the title of the article, author's name (no degrees) author's affiliation, e-mail addresses, mailing address of the corresponding author, and running head less than 60 characters.

ABSTRACT, KEY WORDS, SUBJECT CLASSIFICATION

The manuscript must be accompanied by a brief abstract, no longer than 100-150 words. It should make minimal use of mathematical symbols and displayed formulas. Mathematics Subject Classification (2010) and a list of 4-5 key words must be given.

FIGURES

Figures should be prepared in a digital form suitable for direct reproduction, at resolution of 300 dpi or higher, and in EPS, TIFF or JPEG format.

REFERENCES

Bibliographic references should be listed alphabetically at the end of the article. The authors should consult Mathematical Reviews for the standard abbreviations of journal names.

PROOFS, PAGE CHARGES, OFFPRINTS

The galley proofs will be sent to corresponding author. Late return of the proofs will delay the article to a later issue.

There are no page charges. Authors will receive PDF file of the printed article free of charge. Additional offprints may be ordered from OaM prior to publication.

FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

OaM is published by Publishing House **ELEMENT**, Zagreb, Croatia.

The publication of this journal is supported by Ministry of science, education and sports, Croatia

All correspondence and subscription orders should be addressed to the Editorial Office:

www.ele-math.com

e-mail: oam@ele-math.com

Operators and Matrices
Editorial Office
Menceticeva 2, 10000 Zagreb, Croatia
Fax: +385 1 6008799

A NOTE ON THE STRUCTURE OF MATRIX * -SUBALGEBRAS WITH SCALAR DIAGONALS

G. MACDONALD¹, L. W. MARCOUX¹, M. MASTNAK¹,
M. OMLADIČ² AND H. RADJAVI

(Communicated by E. Poon)

Abstract. We characterize those unital, self-adjoint algebras of complex $n \times n$ matrices that are simultaneously unitarily similar to algebras in which every member has a scalar diagonal.

1. Introduction

This short paper is concerned with the structure of the set of diagonal entries of complex matrices in an algebra, group or semigroup. More specifically, we are interested in characterising those * -subalgebras \mathcal{A} of $M_n(\mathbb{C})$ for which there exists an orthonormal basis with respect to which each element $A \in \mathcal{A}$ admits a diagonal, all of whose entries are equal. We shall say that such an algebra has *scalar diagonals*.

The question “What can we say about the diagonal of a matrix up to unitary similarity?” has a rich history; below we only mention a few highlights. This short note is, as far as we know, the first to explore a related question for collections of matrices. In our setting we restrict ourselves exclusively to the finite-dimensional setting. When we say a matrix, we always mean a matrix of finite dimensions with complex entries.

A well-known fact about *any* single matrix A is that it is unitarily similar to a matrix with scalar diagonal (see, for example, page 109 of [3]); i.e., for some unitary matrix U , the matrix U^*AU has all of its diagonal entries equal, or $\text{Diag}(U^*AU) = \lambda I$ (with $\lambda = \frac{1}{n} \text{tr}(A)$, where n is the size of A). This result also has infinite-dimensional generalizations [1]. If general similarities are allowed, then every sequence (x_1, x_2, \dots, x_n) satisfying $\text{tr}(A) = \sum_{k=1}^n x_k$ occurs as the sequence of diagonal entries in some matrix similar to A . (See, e.g., [5] or [2], where the statement is proved for matrices over an arbitrary field.)

We also mention the interesting characterisation of diagonals of various types of matrices. For example, Horn [4] shows that the real sequence (d_1, d_2, \dots, d_n) is the diagonal of a rotation matrix of order n (i.e., a real orthogonal $n \times n$ matrix of determinant 1) if and only if it is in the convex hull of those sequences $(\pm 1, \pm 1, \dots, \pm 1)$.

Mathematics subject classification (2010): 15A04, 15A21.

Keywords and phrases: Unitary similarity, scalar diagonals, self-adjoint subalgebras of matrices.

¹ Research is supported in part by NSERC (Canada).

² Research is supported in part by the Slovenian Research Agency (research core funding No. P1-0222).

in which -1 occurs an even number of times (possibly zero times). The same paper shows that if $d_i \geq 0$ for all i , then (d_1, d_2, \dots, d_n) is the diagonal of a rotation matrix if and only if it is the diagonal of a doubly stochastic matrix: whence the well-known Schur-Horn Theorem follows. (That theorem states that there exists a complex $n \times n$ hermitian matrix with eigenvalues $\lambda_1 \leq \dots \leq \lambda_n$ and diagonal entries $d_1 \leq \dots \leq d_n$ if and only if for each $i = 1, \dots, n-1$ we have that $d_1 + \dots + d_i \leq \lambda_1 + \dots + \lambda_i$ and $d_1 + \dots + d_n = \lambda_1 + \dots + \lambda_n$. Analogous result also holds for real orthogonal matrices.)

We are interested in collections of matrices with scalar diagonals. Perhaps the simplest example of a (non-trivial, multiplicative) group of matrices with this property is that generated by a cyclic permutation matrix, where the diagonal of each member consists of either zeros or ones. With a little care, one can actually show that if A is any invertible normal matrix, there is a unitary matrix U such that the group generated by U^*AU consists of matrices with scalar diagonals. In fact, it is a special case of our main result that every commutative, self-adjoint group of matrices (or the algebra generated by such a group) has the scalar-diagonal property after a suitable unitary similarity.

We wish to characterise, up to unitary similarity, those self-adjoint subalgebras (equivalently, semigroups) of $\mathbb{M}_n(\mathbb{C})$, every member of which has a scalar diagonal. Such an algebra \mathcal{A} will necessarily have non-trivial invariant subspaces (if $n > 1$). This is a consequence of Burnside's Theorem, which states that the only subalgebra of $\mathbb{M}_n(\mathbb{C})$ without non-trivial invariant subspaces is $\mathbb{M}_n(\mathbb{C})$ itself. Clearly $\mathbb{M}_n(\mathbb{C})$ does not admit scalar diagonals.

One can in fact show that, under the hypothesis, the number of mutually orthogonal invariant subspaces for \mathcal{A} is no less than \sqrt{n} . But our characterisation will imply more: \mathcal{A} is unitarily similar to a direct sum of algebras $\mathbb{M}_{m_i}(\mathbb{C}) \otimes I_{k_i}$, where $m_i \leq k_i$ for each i .

Since a unital, self-adjoint semigroup \mathcal{S} , e.g. a group of unitary matrices, has the scalar-diagonal property if and only if the algebra generated by it does, the main result has an easy corollary for such semigroups.

We take the opportunity to thank Leo Livshits, who graciously organised the meeting that led to this work, but could not attend it himself. We would also thank the unnamed referees for their helpful suggestions.

2. Notation and preliminary observations

We use the convention whereby *we start counting at 0*. In particular, we enumerate rows and columns of an $n \times n$ matrix by the integers $0, 1, \dots, (n-1)$ with the convention that the row and column indices are taken modulo n . For example, if $n = 3$ and

$$A = (a_{i,j})_{i,j=0}^{n-1} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \in \mathcal{M}_3(\mathbb{C}),$$

then $a_{0,0} = a_{3,0} = 1$, $a_{1,2} = a_{7,-1} = 6$, the 0-row of A is $(1\ 2\ 3)$, and the 1-column of A is $(2\ 5\ 8) := \begin{pmatrix} 2 \\ 5 \\ 8 \end{pmatrix}$. For an integer $n \geq 1$, we consider elements of the vector

space \mathbb{C}^n to be column vectors; we often use the usual comma-delimited row notation for them (as at the end of the preceding sentence).

For a strictly positive integer n , we use $\omega_n = e^{2\pi i/n}$ to denote a fixed primitive n -th root of 1 and we use F_n to denote the $n \times n$ discrete Fourier transform (DFT) matrix

$$F_n = \frac{1}{\sqrt{n}} \left(\omega_n^{jk} \right)_{j,k=0}^{n-1} \in \mathcal{M}_n(\mathbb{C}).$$

For example

$$F_1 = (1), \quad F_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad F_3 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega_3 & \omega_3^2 \\ 1 & \omega_3^2 & \omega_3 \end{pmatrix}.$$

It is well known (and easy to see) that F_n is a unitary matrix.

The space of all $n \times n$ matrices with scalar diagonal is denoted by \mathcal{SD}_n , i.e.,

$$\mathcal{SD}_n = \left\{ (a_{j,k})_{j,k=0}^{n-1} \in \mathcal{M}_n(\mathbb{C}) : a_{0,0} = a_{1,1} = \dots = a_{n-1,n-1} \right\} \subseteq \mathcal{M}_n(\mathbb{C}).$$

The space of all $n \times n$ matrices whose non-main-diagonal circulant sums are 0 (we call the sum in the definition below a *circulant sum*) is denoted by \mathcal{CCS}_n , i.e.,

$$\mathcal{CCS}_n = \left\{ (a_{j,k})_{j,k=0}^{n-1} \in \mathcal{M}_n(\mathbb{C}) : \forall m \in \{1, \dots, n-1\}, \sum_{j=0}^{n-1} a_{j,j+m} = 0 \right\} \subseteq \mathcal{M}_n(\mathbb{C}).$$

We use \mathcal{ZDS}_n to denote the space of all matrices whose non-main-diagonal diagonal sums are 0, i.e.,

$$\mathcal{ZDS}_n = \left\{ (a_{j,k})_{j,k=0}^{n-1} : \forall m \in \{1, \dots, n-1\}, \sum_{j=0}^{n-m} a_{j,j+m} = 0 = \sum_{j=0}^{n-m} a_{j+m,j} \right\} \subseteq \mathcal{M}_n(\mathbb{C}).$$

In the proof of the main result of the paper we will use the fact that $\mathcal{ZDS}_n \subseteq \mathcal{CCS}_n$ and the following lemma.

LEMMA 2.1. $F_n(\mathcal{CCS}_n)F_n^* = \mathcal{SD}_n$.

Proof. Since each of the spaces $\mathcal{CCS}_n, F_n(\mathcal{CCS}_n)F_n^*$ and \mathcal{SD}_n has dimension $n^2 - n + 1$, it is sufficient to prove that $F_n(\mathcal{CCS}_n)F_n^* \subseteq \mathcal{SD}_n$. Let $A = (a_{i,j})_{i,j=0}^{n-1} \in \mathcal{CCS}_n$. That is, for all $k \in \{1, \dots, n-1\}$ we have that $\sum_{i=0}^{n-1} a_{i,i+k} = 0$. We now compute $b_{i,i}$, the (i,i) -entry of $B = F_n A F_n^*$. If $\zeta = \omega_n^i$, then (recall that we are taking indices modulo n):

$$\begin{aligned} nb_{i,i} &= (1 \ \zeta \ \dots \ \zeta^{n-1}) A \begin{pmatrix} \frac{1}{\zeta} \\ \vdots \\ \overline{\zeta}^{n-1} \end{pmatrix} \\ &= \left(\sum_{j=0}^{n-1} \zeta^j a_{j,0} \ \sum_{j=0}^{n-1} \zeta^j a_{j,1} \ \dots \ \sum_{j=0}^{n-1} \zeta^j a_{j,n-1} \right) \begin{pmatrix} \frac{1}{\zeta} \\ \vdots \\ \overline{\zeta}^{n-1} \end{pmatrix} \end{aligned}$$

$$\begin{aligned}
&= \sum_{k=0}^{n-1} \sum_{j=0}^{n-1} \zeta^j a_{j,k} \bar{\zeta}^k = \sum_{j=0}^{n-1} \sum_{k=0}^{n-1} \zeta^{j-k} a_{j,k} \\
&\stackrel{r=k-j}{=} \sum_{j=0}^{n-1} \sum_{r=0}^{n-1} \zeta^{-r} a_{j,j+r} = \sum_{r=0}^{n-1} \sum_{j=0}^{n-1} \zeta^{-r} a_{j,j+r} \\
&= \sum_{r=0}^{n-1} \zeta^{-r} \sum_{j=0}^{n-1} a_{j,j+r} = \sum_{j=0}^{n-1} a_{j,j} + \sum_{r=1}^{n-1} \zeta^{-r} \sum_{j=0}^{n-1} a_{j,j+r} \\
&= \sum_{j=0}^{n-1} a_{j,j} + \sum_{r=1}^{n-1} \zeta^{-r} \cdot 0 = \sum_{j=0}^{n-1} a_{j,j} = \text{tr}(A). \quad \square
\end{aligned}$$

Throughout the paper we use the canonical isomorphism $\mathcal{M}_{n_1}(\mathbb{C}) \otimes \mathcal{M}_{n_2}(\mathbb{C}) \xrightarrow{\sim} \mathcal{M}_{n_1 n_2}(\mathbb{C})$ given by identifying $A \otimes B$ with the $n_2 \times n_2$ block matrix whose (i, j) -block is the $n_1 \times n_1$ matrix $b_{i,j}A$. In particular, we identify $A \otimes I_n$ with the block diagonal matrix $\text{diag}(A, \dots, A)$.

We use $\langle -, - \rangle$ to denote the standard inner product on \mathbb{C}^n , i.e., for column vectors $x = (x_0, \dots, x_{n-1})$ and $y = (y_0, \dots, y_{n-1})$, here written in the comma-delimited row notation, we have $\langle x, y \rangle = \sum_{i=0}^{n-1} x_i \bar{y}_i$. If $X \subseteq \mathbb{C}^n$, then $X^\perp = \{y \in \mathbb{C}^n : \forall x \in X, \langle x, y \rangle = 0\}$ denotes its orthogonal complement. We employ the same notations $\langle -, - \rangle$ and $(-)^\perp$ to denote the standard inner product and orthogonal complement on $\mathcal{M}_n(\mathbb{C})$. That is, if $A = (a_{i,j})_{i,j=0}^{n-1}$ and $B = (b_{i,j})_{i,j=0}^{n-1}$, then

$$\langle A, B \rangle = \text{tr}(AB^*) = \sum_{i,j=0}^{n-1} a_{i,j} \bar{b}_{i,j}.$$

If $\mathcal{L} \subseteq \mathcal{M}_n(\mathbb{C})$, then $\mathcal{L}^\perp = \{A \in \mathcal{M}_n(\mathbb{C}) : \forall X \in \mathcal{M}_n(\mathbb{C}), \langle A, X \rangle = 0\}$.

We use e_0, \dots, e_{n-1} to denote the standard basis of \mathbb{C}^n . We use $C_i: \mathcal{M}_n(\mathbb{C}) \rightarrow \mathbb{C}^n$, $i \in \{0, \dots, n-1\}$ to denote the projection to column i , i.e., if $A \in \mathcal{M}_n(\mathbb{C})$, then $C_i(A) = Ae_i$. We use $Q_i: \mathbb{C}^n \rightarrow \mathcal{M}_n(\mathbb{C})$, $i = 0, \dots, n-1$ to denote the injection into the column i , i.e., for a column vector $x \in \mathbb{C}^n$ we have $Q_i(x) = xe_i^*$.

3. Matrix $*$ -algebras that have scalar diagonal

LEMMA 3.1. *Let D be a diagonal matrix of trace 0 and let $\mathcal{A} \subseteq \mathcal{M}_n(\mathbb{C})$ be a unital $*$ -algebra. If $\mathcal{A} \subseteq \mathcal{SD}_n$, then $\mathcal{A}D \subseteq \mathcal{A}^\perp$.*

Proof. Let $A, X \in \mathcal{A}$. Since \mathcal{A} is a $*$ -algebra we have that $A^*X \in \mathcal{A} \subseteq \mathcal{SD}_n$, hence A^*X has a scalar diagonal, and therefore $0 = \text{tr}(D^*(A^*X)) = \langle X, AD \rangle$. \square

LEMMA 3.2. *Let $\mathcal{A} \subseteq \mathcal{M}_n(\mathbb{C})$ be a $*$ -algebra. If $\mathcal{A} \subseteq \mathcal{SD}_n$, then for each $i = 0, \dots, n-1$ we have that $\dim \mathcal{A} = \dim C_i(\mathcal{A})$.*

Proof. We will prove that $\dim \mathcal{A} = \dim C_0(\mathcal{A})$. The claim of the lemma then follows from the fact that the condition $\mathcal{A} \subseteq \mathcal{SD}_n$ is invariant under any permutational

similarity. Indeed, if $\mathcal{A} \subseteq \mathcal{SD}_n$ and P is the permutation matrix corresponding to the transposition that switches 0 and j , then $P\mathcal{A}P \subseteq \mathcal{SD}_n$ and $C_0(P\mathcal{A}P) = P(C_j(\mathcal{A}))$.

For $i = 0, \dots, n-1$ let $\mathcal{C}_i = C_i(\mathcal{A})$, let $k_i = \dim \mathcal{C}_i$, and let $\mathcal{Q}_i = Q_i(\mathcal{C}_i^\perp)$. It is clear that for all i we have that $\mathcal{Q}_i \subseteq \mathcal{A}^\perp$.

Let

$$\mathcal{X} = \text{Span}\{AD : A \in \mathcal{A}, D \text{ diagonal, } \text{tr}(D) = 0\}.$$

By Lemma 3.1 we have that $\mathcal{X} \subseteq \mathcal{A}^\perp$. We claim that $\dim \mathcal{X} \geq \sum_{i=1}^{n-1} k_i$. To this end let, for $i = 1, \dots, n-1$, $\tilde{C}_i : \mathcal{C}_i \rightarrow \mathcal{A}$ denote a fixed right linear inverse of the surjective linear map $C_i|_{\mathcal{A}} : \mathcal{A} \rightarrow \mathcal{C}_i$ (e.g., find C_i -pre-images of a basis of \mathcal{C}_i and extend by linearity). Also note that for $D_i = -e_0e_0^* + e_ie_i^*$, we have that M_{D_i} , the right multiplication by D_i , is equal to $-Q_0C_0 + Q_iC_i$. Using this fact it is easy to see that the linear map

$$F : \mathcal{C}_1 \times \dots \times \mathcal{C}_{n-1} \rightarrow \mathcal{X}$$

given by

$$F(x_1, \dots, x_{n-1}) := \sum_{i=1}^{n-1} M_{D_i}(\tilde{C}_i(x_i)) = (-\sum_{i=1}^{n-1} Q_0C_0\tilde{C}_i(x_i) x_1 \dots x_{n-1}) \in \mathcal{X},$$

is a well-defined linear injection; so $\sum_{i=1}^{n-1} k_i = \dim(\mathcal{C}_1 \times \dots \times \mathcal{C}_{n-1}) \leq \dim \mathcal{X}$.

Since the spaces $\mathcal{X}, \mathcal{Q}_0, \dots, \mathcal{Q}_{n-1}$, are pairwise orthogonal we therefore have that

$$\begin{aligned} \dim \mathcal{A}^\perp &\geq \dim(\mathcal{X} + \mathcal{Q}_0 + \dots + \mathcal{Q}_{n-1}) \\ &= \dim \mathcal{X} + \sum_{i=0}^{n-1} \dim \mathcal{Q}_i \\ &\geq \sum_{i=1}^{n-1} k_i + \sum_{i=0}^{n-1} (n - k_i) = n^2 - k_0. \end{aligned}$$

Hence

$$\dim \mathcal{A} = n^2 - \dim \mathcal{A}^\perp \leq n^2 - (n^2 - k_0) = k_0 = \dim C_0(\mathcal{A}) \leq \dim \mathcal{A},$$

and thus $\dim \mathcal{A} = \dim C_0(\mathcal{A})$. \square

LEMMA 3.3. *Let $\mathcal{A} = \mathcal{M}_m \otimes I_n \oplus 0_p \subseteq \mathcal{M}_{mn+p}$. If \mathcal{A} is unitarily similar to a subset of \mathcal{SD}_{mn+p} , then $m \leq n$.*

Proof. Let U be a unitary matrix such that $U\mathcal{A}U^* \subseteq \mathcal{SD}_{mn+p}$.

Since \mathcal{A} has a common range of dimension mn , we have that $\dim C_0(U\mathcal{A}U^*) \leq mn$. In more detail: let $\mathcal{X} \subseteq \mathbb{C}^{mn+p}$ be the subspace spanned by the first mn basis vectors e_0, \dots, e_{mn-1} . Then for each $f \in \mathbb{C}^{mn+p}$, the space $\mathcal{A}f$ is contained in \mathcal{X} . Therefore $C_0(U\mathcal{A}U^*) = U(\mathcal{A}(U^*e_0))$ is contained in $U(\mathcal{X})$.

Hence by Lemma 3.2 we have that

$$m^2 = \dim \mathcal{A} = \dim(U\mathcal{A}U^*) = \dim C_0(U\mathcal{A}U^*) \leq mn,$$

and therefore $m \leq n$. \square

LEMMA 3.4. *Let $\mathcal{A} = \mathcal{M}_m(\mathbb{C}) \otimes I_n \subseteq \mathcal{M}_{mn}(\mathbb{C})$ with $m \leq n$. Let $D_{m,n} = \text{diag}(1, \omega_n, \dots, \omega_n^{m-1}) \in \mathcal{M}_m(\mathbb{C})$ and let $\tilde{D}_{m,n} = \text{diag}(I_m, D_{m,n}, \dots, D_{m,n}^{n-1}) \in \mathcal{M}_{mn}$. Then $\tilde{D}_{m,n} \mathcal{A} \tilde{D}_{m,n}^* \subseteq \mathcal{ZDS}_{mn}$.*

Proof. Let $A = (a_{i,j})_{i,j=0}^{m-1} \in \mathcal{M}_m(\mathbb{C})$, let $B = (b_{i,j})_{i,j=0}^{mn-1} = \tilde{D}_{m,n}(A \otimes I_n)\tilde{D}_{m,n}^*$. We will prove that all strictly upper diagonal sums $\sum_{i=0}^{mn-1-k} b_{i,i+k}$, $k \in \{1, \dots, mn-1\}$ are zero (the proof of the fact that all strictly lower diagonal sums are zero is symmetric). Let $k \in \{1, \dots, mn-1\}$. Note that for $p \in \{0, \dots, m-1\}$, and $q \in \{0, \dots, n-1\}$, we have that

$$b_{mq+p, mq+p+k} = \begin{cases} \omega_n^{-kq} a_{p,p+k} & \text{if } k \leq m-1-p \\ 0 & \text{if } k > m-1-p \end{cases}.$$

Hence for $k > m-1-p$ we clearly have that $\sum_{i=0}^{mn-1-k} b_{i,i+k} = 0$. Now assume that $k \leq m-1-p$. Since $m \leq n$ we have $\omega_n^k \neq 1$ and thus $\sum_{q=0}^{n-1} \omega_n^{-kq} = 0$. Therefore

$$\begin{aligned} \sum_{i=0}^{mn-1-k} b_{i,i+k} &= \sum_{p=0}^{m-1-k} \sum_{q=0}^{n-1} b_{mq+p, mq+p+k} \\ &= \sum_{p=0}^{m-1-k} \sum_{q=0}^{n-1} \omega_n^{-kq} a_{p,p+k} \\ &= \sum_{p=0}^{m-1-k} a_{p,p+k} \sum_{q=0}^{n-1} \omega_n^{-kq} = 0. \quad \square \end{aligned}$$

THEOREM 3.5. *Let \mathcal{A} be a unital $*$ -subalgebra of $\mathcal{M}_n(\mathbb{C})$. Then \mathcal{A} is unitarily similar to a subspace of \mathcal{SD}_n if and only if up to unitary similarity we have that $\mathcal{A} = \bigoplus_{i=1}^k \mathcal{M}_{m_i} \otimes I_{n_i}$ with $m_i \leq n_i$ for $i = 1, \dots, k$.*

Proof. We assume, with no loss of generality, that $\mathcal{A} = \bigoplus_{i=1}^k \mathcal{M}_{m_i} \otimes I_{n_i}$ for some natural numbers k , m_1, \dots, m_k , n_1, \dots, n_k .

(\implies): Assume that \mathcal{A} is unitarily similar to a subspace of \mathcal{SD}_n . For each $i \in \{1, \dots, k\}$, \mathcal{A} has a subalgebra \mathcal{A}_i unitarily similar to $\mathcal{M}_{m_i} \otimes I_{n_i} \oplus 0_{n-m_i n_i}$. Since each \mathcal{A}_i is also unitarily similar to a subspace of \mathcal{SD}_n we have, by Lemma 3.3, that $m_i \leq n_i$ for all i .

(\impliedby): Assume that for all $i = 1, \dots, k$ we have that $m_i \leq n_i$. Let $D = \text{diag}(\tilde{D}_{m_1, n_1}, \dots, \tilde{D}_{m_k, n_k})$. By Lemma 3.4 we have that $D \mathcal{A} D^* \subseteq \mathcal{ZDS}_n$. Since $\mathcal{ZDS}_n \subseteq \mathcal{ZCS}_n$ we therefore have, by Lemma 2.1, that $F_n D \mathcal{A} D^* F_n^* \subseteq \mathcal{SD}_n$. \square

COROLLARY 3.6. *Every unital, commutative, self-adjoint semigroup of matrices – in particular, every commutative group of unitaries – is unitarily similar to a semi-group of matrices with scalar diagonal.*

Proof. Let \mathcal{S} be a unital, commutative, self-adjoint semigroup of matrices. We can assume with no loss of generality that \mathcal{S} consists of diagonal matrices. Then so

does the algebra \mathcal{A} generated by \mathcal{S} . Just note that \mathcal{A} is a direct sum of algebras of the form $\mathbb{M}_{m_i} \otimes I_{n_i}$ with $m_i = 1$ for all i . \square

The corollary above also follows from Lemma 2.1 which gives an explicit formula for the unitary similarity in question.

REFERENCES

- [1] P. FAN, *On the diagonal of an operator*, Trans. Amer. Math. Soc. 283, no. 1: 239–251, 1984.
- [2] P. A. FILLMORE, *On similarity and the diagonal of a matrix*, Amer. Math. Monthly 76: 167–169, 1969.
- [3] P. R. HALMOS, *Finite Dimensional Vector Spaces*, Van Nostrand, Princeton, N.J., 1958.
- [4] A. HORN, *Doubly stochastic matrices and the diagonal of a rotation*, Amer. J. Math. 76: 620–630, 1954.
- [5] L. MIRSKY, *Matrices with prescribed characteristic roots and diagonal elements*, J. London Math. Soc. 33: 14–21, 1958.

(Received October 1, 2019)

G. MacDonald

School of Mathematical and Computational Sciences
University of Prince Edward Island
Charlottetown, PE, Canada C1A 4P3
e-mail: gmacdonald@upei.ca

L. W. Marcoux

Department of Pure Mathematics
University of Waterloo
Waterloo, ON, Canada N2L 3G1
e-mail: LWMarcoux@uwaterloo.ca

M. Mastnak

Department of Mathematics and Computing Science
Saint Mary's University
923 Robie St, Halifax, Nova Scotia, Canada B3N 1Z9
e-mail: mmastnak@cs.smu.ca

M. Omladič

Institute of Mathematics, Physics and Mechanics
Jadranska 19, 1000 Ljubljana, Slovenia
e-mail: matjaz@omladic.net

H. Radjavi

Department of Pure Mathematics
University of Waterloo
Waterloo, ON, Canada N2L 3G1
e-mail: hradjavi@uwaterloo.ca

